



Ranunculus Millani ;

FERD. VON MUELLER,

Transactions of the Philosophical Society of Victoria 1855, pag. 97.

ORDER RANUNCULACEÆ.

Alps of Victoria and New South Wales ; Mount Kosciusko,
5500—6500 feet high.

Contributor : Baron von Mueller.



Hibbertia angustifolia;

SALISBURY,

Paradisus Londinensis, pag. 73. (1806.)

ORDER DILLENIACEÆ.

Victoria and Tasmania.

Port Phillip.—Contributor : G. Luehmann.



Bursaria spinosa ;

CAVANILLES,

Icones et Descriptiones Plantarum, vol. iv. pag. 30, tab. 350. (1797.)

ORDER PITTOSPOREÆ.

Through the greater part of the Australian Continent.

Port Phillip.—Contributor : C. Groener.



Drosera Whittackeri ;

PLANCHON,

Annales des Sciences Naturelles, troisième série, vol. ix. pag. 302. (1848.)

ORDER DROSERACEÆ.

Victoria and South Australia.

Port Phillip.—Contributor : C. Richards.



Tetratheca ciliata ;

LINDLEY,

Sir Th. Mitchell's Three Expeditions into Eastern Australia, vol. ii. pag. 206.
(1838.)

ORDER TREMANDREÆ.

From the eastern parts of South Australia through the colony
Victoria into New South Wales, also in Tasmania.

Port Phillip.—Contributor : C. Walter.



Lasiopetalum Baueri ;

STEETZ,

Lehmann Plantæ Preissianæ, vol. ii. pag. 339. (1847.)

ORDER STERCULIACEÆ.

South Australia, Victoria, and New South Wales.

Port Phillip.—Contributor : Baron von Mueller.



Correa speciosa ;

W. T. AITON,
Epitome of the Hortus Kewensis, pag. 366. (1814.)

ORDER RUTACEÆ.

From the Great Bight to South Queensland; also in Tasmania.

Port Phillip.—Contributor : C. Richards.



Eriostemon squameus ;

LA BILLARDIÈRE,

Novæ Hollandiæ Plantarum Specimen, vol. i. pag. 111, tab. 141. (1804.)

New South Wales, Victoria, Tasmania.

Cape Otway.—Contributor : Baron von Mueller.



Pultenæa rosea ;

FERD. VON MUELLER,

Fragmenta Phytographiæ Australiæ, vol. ii. pag. 15. (1860.)

ORDER LEGUMINOSÆ.

Victoria ; Mount William, 4000—5000 feet high.

Contributor : D. Sullivan.



Acacia longifolia ;

WILLDENOW,
Species Plantarum, vol. iv. pag. 1052. (1805.)

ORDER LEGUMINOSÆ.

From South Australia to South Queensland.
Gippsland.—Contributor : Baron von Mueller.



Lhotzkya genethylloides ;

FERD. VON MUELLER,

Transactions of the Philosophical Society of Victoria, vol. i. pag. 16. (1854.)

ORDER MYRTACEÆ.

Eastern part of South Australia and Western part of Victoria.

Grampians.—Contributor : D. Sullivan.



Kunzea Muelleri ;


BENTHAM,

Flora Australiensis, vol. iii. pag. 113. (1866.)

ORDER MYRTACEÆ.

Alps of Victoria and New South Wales ; Mount Kosciusko,
5000—7000 feet high.

Contributor : Baron von Mueller.



Eucalyptus viminalis ;

LA BILLARDIÈRE,

Novæ Hollandiæ Plantarum Specimen, vol. ii. pag. 12, tab. 151. (1806.)

ORDER MYRTACEÆ.

South Australia, Victoria, Tasmania, and New South Wales.

Port Phillip.—Contributor : C. Groener.



Ricinocarpus pinifolius ;


DESFONTAINES,

Memoires du Muséum d'Histoire Naturelle, Paris, vol. iii. pag. 459, tab. 22. (1818.)

ORDER EUPHORBIACEÆ.

Victoria, Tasmania, New South Wales, South Queensland.

Port Phillip.—Contributor : C. Walter.



Amperea spartioides ;

BROGNIART,

Duperrey Voyage de la Coquille, Botanique, pag. 226, tab. 49. (1829.)

ORDER EUPHORBIACEÆ.

South Australia, Victoria, Tasmania, New South Wales.

Port Phillip.—Contributor : G. Luehmann.



Pomaderris elliptica ;

LA BILLARDIÈRE,

Novæ Hollandiæ Plantarum Specimen, vol. i. pag. 61, tab. 86. (1804.)

ORDER RHAMNACEÆ.

Victoria, Tasmania, New South Wales.

Dandenong.—Contributor : G. Luehmann.



Spyridium parvifolium ;

FERD. VON MUELLER,
Fragmenta Phytographiæ Australiae, vol. iii. pag. 79. (1862.)

ORDER RHAMNACEÆ.

South Australia, Victoria, Tasmania, New South Wales.

Dandenong.—Contributor : G. Luehmann.



Bauera sessiliflora ;

FERD. VON MUELLER,

Transactions of the Philosophical Society of Victoria 1855, pag. 41.

ORDER SAXIFRAGEÆ.

Victoria.

Mount William.—Contributor : D. Sullivan.



Tillæa macrantha ;

J. HOOKER,
Hooker, Icones Plantarum, 310. (1840.)

ORDER CRASSULACEÆ.

Through a large extent of extratropical Australia.

Port Phillip.—Contributor : C. Groener.



Haloragis micrantha ;

R. BROWN,

Appendix to Flinders' Voyage, pag. 550. (1814.)

ORDER HALORAGÆ.

From South Australia to South Queensland ; also in Tasmnia.

Port Phillip.—Contributor : C. Groener.




Hydrocotyle geraniifolia ;

FERD. VON MUELLER,
Transactions of the Victorian Institute 1855, pag. 126.

ORDER UMBELLIFERÆ.

Victoria, New South Wales.

Upper Yarra.—Contributor : Baron von Mueller.



Trachymene australis ;

BENTHAM,

Flora Australiensis, vol. iii. pag. 349. (1866.)

ORDER UMBELLIFERÆ.

Extra-tropical and eastern tropical Australia.

Port Phillip.—Contributor : G. Luehmann.



Aster stellulatus;

LA BILLARDIÈRE,
Novæ Hollandiæ Plantarum Specimen, vol. ii. pag. 50, tab. 196. (1806.)

ORDER COMPOSITÆ.

Victoria, Tasmania, New South Wales, Queensland.

Dandenong.—Contributor : G. Luehmann.



Helichrysum apiculatum;

CANDOLLE,

Prodromus Systematis Naturalis Regni Vegetabilis, vol. vi. pag. 195. (1837.)

ORDER COMPOSITÆ.

Nearly over all Australia.

Port Phillip.—Contributor : G. Luehmann.



Leptorrhynchus squamatus ;

LESSING,

Synopsis Generum Compositarum, pag. 273. (1832.)

ORDER COMPOSITÆ.

South Australia, Victoria, New South Wales, Tasmania.

Port Phillip.—Contributor : C. Groener.



Wahlenbergia gracilis ;

A. DE CANDOLLE,
Monographie des Campanulées, pag. 142. (1830.)

Throughout Australia.

Port Phillip.—Contributor : C. Groener.



Brunonia australis ;

SMITH,

Transactions of the Linnean Society, vol. x. pag. 367, tab. 28. (1808.)

ORDER GOODENOVIÆ.

Through the greater part of extra-tropical Australia.

Port Phillip.—Contributor : C. Groener.



Epacris impressa ;

LA BILLARDIÈRE,

Novæ Hollandiæ Plantarum Specimen, vol. i. pag. 43, tab. 58. (1804.)

ORDER EPACRIDEÆ.

South Australia, Victoria, Tasmania, New South Wales.

Port Phillip.—Contributor : G. Luehmann.



Styphelia virgata ;

LA BILLARDIÈRE,

Novæ Hollandiæ Plantarum Specimen, vol. i. pag. 46, tab. 64. (1804.)

Leucopogon virgatus, R. Brown, Prodrômus Floræ Novæ
Hollandiæ, pag. 543. (1810.)

ORDER EPACRIDEÆ.

Extensively spread through South-East Australia, penetrating to
the Alps and deserts.

Port Phillip.—Contributor : C. Walter.



Poirertia cucullata ;

CAVANILLES,

Icones et Descriptiones Plantarum, tab. 343. (1797.)

Sprengelia incarnata, Smith, Tracts relating to Natural History,
pag. 272, tab. 2. (1798.)

ORDER EPACRIDEÆ.

From South Australia through the colony Victoria to New South
Wales, also in Tasmania, ascending to alpine regions.

Port Phillip.—Contributor : C. Walter.



Richea procera ;

FERD. VON MUELLER,
Fragmenta Phytographiæ Australiæ, vol. vi, pag. 68.

ORDER EPACRIDEÆ.

Tasmania.

Mount Wellington.—Contributor : Samuel Hannafor.



Erithræa australis ;

R. BROWN,

Prodromus Floræ Novæ Hollandiæ, pag. 451. (1810.)

ORDER GENTIANEÆ.

In most parts of Australia.

Port Phillip.—Contributor : G. Luehmann.



Euphrasia Brownii ;

FERD. VON MUELLER,
Fragmenta Phytographiæ Australiæ, vol. v. pag. 88. (1865.)

ORDER SCROPHULARINÆ.

Through extra-tropical Australia.

Dandenong.—Contributor : C. Walter.



Prostanthera nivea ;

CUNNINGHAM,

Bentham Labiatarum Genera et Species, pag. 452. (1835.)

ORDER LABIATÆ.

Victoria and New South Wales.

You Yangs.—Contributor : C. Walter.



Grevillea alpina ;

LINDLEY,

Mitchell's Three Expeditions, vol. ii. pag. 179. (1838.)

ORDER PROTEACEÆ.

Mount Disappointment.

Contributor : C. Walter.



Pimelea axiflora ;

FERD. VON MUELLER,
First General Report, pag. 17. (1853.)

ORDER THYMELEÆ.

Victoria and New South Wales.

Upper Yarra.—Contributor : C. Walter.



Thelymitra antennifer ;

J. HOOKER,

Flora Tasmanica, vol. ii. pag. 4 ; plat. 101, fig. A. (1860.)

ORDER ORCHIDÆ.

From the southern parts of New South Wales to West Australia
and Tasmania.

Port Phillip.—Contributor : G. Luehmann.



Cyrtostylis reniformis ;

R. BROWN,

Prodromus Floræ Novæ Hollandiæ, pag. 322. (1810.)

ORDER ORCHIDÆ.

From West Australia to Queensland, particularly in coast regions.

Port Phillip.—Contributor : C. Walter.



Hypoxis glabella ;

R. BROWN,

Prodromus Floræ Novæ Hollandiæ, pag. 289. (1810.)

ORDER AMARYLLIDEÆ.

South Australia, Victoria, Tasmania.

Port Phillip.—Contributor : C. Groener.



Anguillaria australis ;

FERD. VON MUELLER,

Fragmenta Phytographiæ Australiæ, vol. vii. pag. 74. (1870.)

VARIETY DIOICA.

Anguillaria dioica, R. Brown, Prodrômus Floræ Novæ Hollandiæ,
pag. 273. (1810.)

ORDER LILIACEÆ.

Widely dispersed through extratropical and eastern tropical
Australia.

Port Phillip.—Contributor : C. Groener.



Burchardia umbellata;

R. BROWN,
Prodromus Floræ Novæ Hollandiæ, pag. 273. (1810.)

ORDER LILIACEÆ.

Nearly throughout extra-tropical Australia.

Port Phillip.—Contributor : C. Groener.



Chamæscilla corymbosa ;

FERD. VON MUELLER,

Fragmenta Phytographiæ Australiæ, vol. vii. pag. 68. (1870.)

Cæsia corymbosa, R. Brown, Prodromus Floræ Novæ Hollandiæ,
pag. 277. (1810.)

ORDER LILIACEÆ.

From New South Wales to Tasmania and West Australia.

Port Phillip.—Contributor : C. Groener.



Hypolaena fastigiata ;

R. BROWN,

Prodromus Floræ Novæ Hollandiæ, pag. 251. (1810.)

ORDER RESTIACEÆ.

South Australia, Victoria, Tasmania, New South Wales.

Port Phillip.—Contributor : G. Luehmann.



Stipa semibarbata ;

R. BROWN,

Prodromus Floræ Novæ Hollandiæ, pag. 174. (1810.)

ORDER GRAMINEÆ.

Southern tracts of Australia.

Port Phillip.—Contributor : G. Luehmann.



Ehrharta stipoides ;


LA BILLARDIÈRE,

Novæ Hollandiæ Plantarum Specimen, vol. i. pag. 191, tab. 118. (1804.)

ORDER GRAMINEÆ.

Through the greater part of extra-tropical Australia.

Yarra.—Contributor : G. Luehmann.



Gleichenia circinata ;

SWARTZ,

Synopsis Filicum, pag. 165 et 394. (1806.)

ORDER FILICES.

South Australia to Queensland, in deep forest valleys.

Upper Yarra.—Contributor : C. Walter.



Tmesipteris Tannensis ;

BERNHARDI,

Schrader's Journal fuer Botanik 1800, sect. ii. pag. 131, tab. 2, fig. 5.

ORDER LYCOPODIACEÆ.

Victoria, Tasmania, New South Wales, Queensland.

Upper Yarra.—Contributor : G. Luehmann.



Dawsonia superba;

GREVILLE,

Annals of Natural History 1847, pag. 226. tab. 12.

ORDER MUSCI.

Victoria, Tasmania, New Zealand, Sources of the Yarra Yarra.

Contributor : Baron von Mueller.